

Health care deprivation profiles in the measurement of inequality and inequity: an application to GP fundholding in the English NHS

Mauro Laudicella^α, Richard Cookson^β, Andrew M. Jones^γ, Nigel Rice^δ
University of York

Working paper for the HESG meeting, 27-29 August 2008

Original working paper downloadable from: <http://www.york.ac.uk/res/herc/research/hedg/wp.htm>

Abstract: This paper proposes a new approach to the measurement of inequality and inequity in the delivery of health care based on contributions from the literature on poverty and deprivation. This approach has some appealing characteristics: 1) inequity is additively decomposable by population subgroups; 2) the approach does not rely on socio-economic ranks; 3) it provides a graphical representation of the distribution of inequity; 4) it offers a range of indices consistent with dominance. An empirical application is provided investigating the effect of the GP fundholding reform on equity in English NHS. The results show that the most equitable GP practices self-selected into the scheme in 1991; evidence of an inequity-reducing treatment effect as well as a self-selection effect are found in 1992 and 1993; the self-selection process reduces and no evidence of a treatment effect is present thereafter.

Keywords: inequality, inequity, health care, poverty, deprivation, dominance, GP fundholding.

JEL classification: D63, I18, I32.

Acknowledgements: Dataset assembly for this project was funded by the ESRC Public Services Programme Phase II Award RES-166-25-0055. Hospital episode statistics were provided by the NHS Health and Social Care Information Centre, on license from the Department of Health. Census data were provided by the MIMAS service at the University of Manchester. Census data for 1991 is licensed by the Office of Population Censuses and Surveys, and census data for 2001 is licensed by the Office of National Statistics.

^α Centre for Health Economics, University of York. Correspondence: Centre for Health Economics, Alcuin "A" Block, University of York, York, YO10 5DD, UK. Tel: +44 1904 321 423. E-mail: ml529@york.ac.uk.

^β Department of Social Policy and Social Work, University of York.

^γ Department of Economics and Related Studies, University of York.

^δ Centre for Health Economics, University of York.

1. Introduction

Horizontal equity in health care is commonly defined as equal treatment for equal need. A large part of the empirical literature in this field focuses only on violations of the principle of horizontal equity that are related to income or socio-economic status and measures inequity using standardized concentration curves and indices (Wagstaff and van Doorslaer, 2000a). These studies fail in providing a measure of inequity associated with variables that do not vary with income or socio-economic status since their tools of analysis do not capture such aspects.

There is evidence that in OECD countries many other variables drive inequity beside income, especially in countries where health care is provided mainly free of charge. Income does not explain the probability of visiting a general practitioner across most of the OECD countries, while it is associated with inequity in visits to a specialist in almost all countries (van Doorslaer et al., 2006). However, income is not always the most important factor in explaining subsequent visits to a specialist (van Doorslaer et al., 2004). Moreover, some authors find evidence that other variables such as ethnicity, education and employment status drive inequity in both primary and secondary care in the English NHS (Morris et al., 2005).

Many of the empirical studies on inequity in health care adopt a regression-based approach (Gravelle et al., 2006). This strategy includes characteristics that are not related to income in the analysis of inequity. Most of these studies model the probability of receiving health care as a function of a set of need variables and a set of other individual characteristics and policy variables. Inequity is found whenever the probability of receiving a treatment depends on variables that do not belong to the set of the need variables. This approach is effective in identifying the presence of inequity as well as the factors driving inequity, but has considerable limitations in quantifying inequity and in making inequity comparisons.

The methodological contribution of this paper is to apply some developments in the literature on poverty and deprivation to the analysis of inequity in health care (Jenkins and Lambert, 1997; Shorrocks, 1998). This produces a new approach to the measurement of inequity that enlarges the scope of the analysis and offers advantages with respect to the previous empirical literature. It provides inequity ranks that are consistent with a wide range of social preferences for equity and offers a useful graphical representation of important aspects of the distribution of inequity in the population. Further, a class of indices consistent with dominance is introduced in order to aggregate the distribution of inequity into a single value.

The new approach allows a decomposition of inequity into population subgroups. Inequity measured in the population as a whole can be decomposed into a weighted sum of inequity experienced in any partition of the population. This property permits inequity rankings across specific demographic groups within the population (e.g. education, or region, or ethnic groups) and identification of those which are particularly suffering inequity. Also, it is possible to analyse variation in inequity associated with discrete variations in any characteristics of interest in the groups considered (e.g. level of education, or age, or travel distance from hospital, or socio-economic status). Thus, the additive decomposability property enlarges substantially the scope of the inequity analysis with respect to previous studies based on concentration curves and indices.

Finally, the measure of inequity does not rely on the socio-economic status (SES) of our observations. We are able to obtain a measure of inequity whenever such information is missing or inaccurate (e.g. when SES is a discrete or categorical variable). Also, we are able to analyse sources of inequity which are unrelated or weakly related to income, such as regional disparities in the supply of care, education, ethnicity, age, and gender.

The second contribution of this paper is an empirical application of the new method to the analysis of the GP fundholding reform in the English NHS. We measure the effect of the reform on equity in the utilization of secondary care for elective hip replacements across the electoral wards of England. GPs act as gatekeepers to secondary care in the English NHS and traditionally bore no financial cost in referring patients to secondary care. In allowing GP practices to hold budgets for hospital care, and to re-invest any savings within the practice, the fundholding reform aimed to give a financial incentive to manage access to secondary care to those GPs who joined the scheme. The consequences of this reform for overall access to secondary care have been intensively studied. There is evidence that GP fundholding practices reduced overall elective admission rates (Dusheiko et al., 2006) while at the same time reducing the average waiting time for secondary care admissions (Propper et al. 2002). However, there is little evidence on equity. We provide measures of the impact of the GP fundholding reform on equity that are consistent with a wide range of social preferences for equity. We use elective hip replacements as an indicator of health care inequity and examine inequity in the distribution of elective hip replacements within the most socio-economically deprived quartiles of electoral wards of England. Hip replacement is performed routinely and is a common, effective and low risk surgical procedure. It is often considered one of the most important examples of health care inequality in small area administrative studies, since it shows a clear socio-economic gradient after controlling for need (Chaturvedi and Ben-Shlomo 1995; Dixon et al., 2004; Cookson, et al., 2007).

2. Health care deprivation profiles as a measure of inequity

This section provides the theoretical framework underlying our approach to the measurement of inequity in the delivery of health care. Its foundations rely on previous work by Jenkins and Lambert (1997) and Shorrocks (1998) in the literature on poverty and deprivation.

Consider a vector of observations on health care utilization: $h = (h_1, h_2, \dots, h_n)$. The elements of the vector h might be number of visits to a specialist, or measure of a specific procedure such as number of elective hip replacements, within a given time period.

Define the vector $h^* = (h^*_1, h^*_2, \dots, h^*_n)$ that for each observation in “ h ” gives its need-expected level of utilization as a function of a set of need indicators “ n ” (e.g. age, sex, self assessed health status, presence of limiting long term illness, etc.):

$$h^* = f(n) \tag{1}$$

This approach does not depend on a specific “need adjustment”. For instance, h^* can be exogenously set by a policy maker as the minimum amount of health care resources that individuals or communities should receive given their need. In this paper, we follow the tradition of the literature on inequity in health care and define h^* as a function of a set of need indicators and observed health care utilization:

$$h^* = f(n; h) \tag{2}$$

Equation (1) and (2) can be specified at either the individual or small area level. When the units of analysis are individuals, the vector h^* in Equation (2) could be computed using the indirect standardization approach described in Wagstaff and van Doorslaer (2000a). Here, we describe the computation of h^* when the units of analysis are small area observations. This should facilitate the illustration of the empirical application in section 4, in which we use electoral wards as unit of analysis. However, the methods that we describe in Section 2 and 3 also apply at the individual level.

$$h_i^* = \sum_{j=1}^J p_{ji} h_j \quad (3)$$

$$h_j = \frac{\sum_{i=1}^n h_{ji}}{\sum_{i=1}^n p_{ji}} \quad (4)$$

Where p_{ji} is the number of people in area i that belong to need class j (e.g. number of males aged 45-54 suffering from arthritis in electoral ward i); h_{ji} is the observed health care utilization in area i for need class j . The value h_i^* can be interpreted as the amount of health care utilization that we would expect to observe in the area i given its population, if each of its need groups had the same utilization rate as the population as a whole (i.e. the national utilization rate h_j). The computation of a need-expected distribution similar to h^* is a well establish approach in many empirical studies on the measurement of inequity (Wagstaff and van Doorslaer, 2000b). Also, in the epidemiology literature h^* defines the number of expected events in the computation of the indirectly standardized incidence rate (where j usually are based on age and sex variables; Breslow and Day, 1987; Clayton and Hills, 1993).

Now, define the vector of health care gaps (HCG), $x = (x_1, x_2, \dots, x_n)$, as the vector of distances between need-expected and actual utilization, when the gap is positive, and zero otherwise:


$$x_i = \max(h_i^* - h_i; 0) \quad (5)$$

Define the health care deprivation profiles¹ (HCD) as:

$$HCD(x; p) = \int_{F^{-1}(1-p)}^{\xi} x dF(x) \quad \forall p \in [0, 1] \quad (6)$$

Where $F(x)$ is the cumulative distribution function of x . For each $p \in [0, 1]$ the HCD profile is the cumulative sum of the first $100 \cdot p$ per cent of HCGs ranked from the largest (ξ) to the smallest (0).

Figure 1. The Heath Care Deprivation Profile.



¹ The HCD profiles can be considered a special case of the deprivation profiles illustrated in Shorrocks (1998).

The HCD profile $HCD(x; p)$ for x is described by a concave curve that rises continuously from the largest to the smallest HCG; the curve becomes flat in the interval $[r, 1]$ where the HCGs equal zero. The HCD profile provides a graphical representation of useful information on incidence, intensity and inequality in the distribution of HCGs across the areas. The length of the interval $[0, r]$ represents the proportion of areas receiving less health care than need expected, i.e. the incidence of areas having positive HCGs:

$$r = F(x = 0) \quad (7)$$

The height of the curve, μ , is equal to the average HCG in the *whole* population of areas, i.e. the intensity of the gap between need expected and observed health care in the population of areas:

$$\mu = \int_0^{\xi} xF(x) \quad (8)$$

The degree of concavity of the HCD is determined by the rate at which HCGs decrease in the areas. This offers a representation of inequality in the distribution of the HCGs: if the HCGs are constant across all those areas receiving less health care than needed, then the curve become a straight line; the more the HCGs are unequally distributed across the areas, the greater is degree of concavity of the curve. When all the areas are not below their need-expected value h_i^* , then all HCGs equal zero and the curve lies on the x-axis.

If we define deprivation in health care utilization as receiving less health care than needed, then the HCD profiles offer a graphical representation of the distribution of deprivation in health care across the areas. Moreover, the HCD profiles provide a measure of horizontal inequity in health care for all the areas below the frontier of need expected utilization h^* . We illustrate this argument through three propositions.

Proposition 1². No areas are deprived in health care if and only if the observed health care utilization equals the need expected utilization in all areas:

$$x_i \equiv \max(h_i^* - h_i; 0) = 0, \forall i \in N \Leftrightarrow h_i = h_i^*, \forall i \in N. \quad (9)$$

Proposition 1 holds for both “health care deprived” and “health care non-deprived areas”, i.e. those areas below and above their need expected health care utilization h^* . The HCG vector rules out the possibility of having all the areas above or below the need expected frontier of health care utilization (i.e. $h_i > h_i^* \forall i$, or $h_i < h_i^* \forall i$). If some area increases its health care consumption, this will result either in subtracting resources from other areas and/or in raising the need expected health care utilization for all. Intuitively, Proposition 1 is a direct consequence of the definition of the vector h^* as a function of the average level of health care utilization observed in the total population (Equations (3) and (4)). Thus, the need expected frontier of health utilization is endogenously determined.

Proposition 2². No areas are deprived in health care if and only if the share of health care utilization equals the need adjusted share of population in all the areas:

² The Proof of proposition 1, 2 and 3 is in the original version of the paper: <http://www.york.ac.uk/res/herc/research/hedg/wp.htm>

$$x_i \equiv \max(h_i^* - h_i; 0) = 0, \forall i \in N \Leftrightarrow \sum_j \left(\frac{p_{ij} h_j}{p_j H} \right) = \frac{h_i}{H}, \forall i \in N \quad (10)$$

p_{ij}/p_j is the share of the total population in the need class j living in the area i ; h_j/H is the share of health care utilization of all the people in the need class j with respect to the population as a whole; h_{ij}/H is the share of utilization of people in the need class j living in the area i . The need adjusted share of the population is obtained from indirect standardization of the population of every area with respect to the J need classes (Equation (3) and (4)). The right hand side of Equation (10) contains a condition of horizontal equity:

$$\text{If } \sum_j \frac{p_{ij} h_j}{p_j H} = \sum_j \frac{p_{i'j} h_j}{p_j H}, \forall i, i' \in N \Rightarrow \frac{h_i}{H} = \frac{h_{i'}}{H}, \forall i, i' \in N \quad (11)$$

If two areas have an equal need adjusted population share, then they should have an equal share of health care utilization. This condition is similar to the one proposed in Wagstaff et al. (1991) with respect to population income groups, and widely accepted in the literature. Proposition 2 states an unequivocal relationship between the condition of *the absence of deprivation* and the condition of *horizontal equity* in the whole population of areas. Conversely, the relationship between *the presence of deprivation* and *horizontal inequity* is weaker:

Proposition 3²: *If an area is deprived in health care, then it is also unequally treated with respect to all other areas:*

$$x_i \equiv \max(h_i^* - h_i; 0) > 0 \Rightarrow \sum_j \left(\frac{p_{ij} h_j}{p_j H} \right) \neq \frac{h_i}{H} \quad (12)$$

By Proposition 3, an area can be unequally treated and not necessarily be deprived in health care. Specifically, all the areas above the frontier of need expected utilization are not deprived in health care and may well have unequal share of health care for equal need. However, if any area is above the frontier of need expected utilization, then at least one area will be deprived in health care by Proposition 1, and the whole population will suffer inequity by Proposition 2 and 3. Thus, the presence of areas above the frontier of health care utilization implies inequity for the population as a whole.

Now, we are able to interpret the HCD profile $HCD(x; p)$ as a measure of both deprivation and inequity in health care:

- i. $HCD(x; p)$ is a continuous, concave and monotonically increasing function defined over x .
- ii. $HCD(x; p)$ lies on the horizontal-axis if and only if all the HCGs are zero; thus if and only if there is no deprivation and no inequity *in the whole population of areas*.
- iii. $HCD(x; p)$ is monotonically increasing in the average HCG μ .
- iv. $HCD(x; p)$ is monotonically increasing in the variance of the distribution of the HCGs.

Properties (i) and (ii) derive directly from the definition of $HCD(x; p)$; (iii) and (iv) are implied by condition (i).

The HCD profile offers a measure of deprivation in the utilization of health care in the whole population of areas, defined as receiving less health care than needed. The reliability of this measure depends on the extent to which the need expected health care utilization h^* is a correct estimator of the population need for health care. h^* is defined as a function of the average utilization in the population as a whole (Equation (3)). Thus, it is assumed that observed and needed health care on average are equal in the total population. Specifically, the HCD profile assumes that on average there is not an over or under utilization of health care in the population with respect to each of the J need classes considered. If a large part of the demand for some health care services is not met by the supply, this assumption may not hold. In this case, the measure of health care deprivation captured by the HCD profiles will be under-estimated.

The HCD profile also provides a measure of horizontal inequity for those areas deprived in health care relatively to the population as whole. Proposition 3 allows an interpretation of the distance between h^* and h as a measure of both deprivation and horizontal inequity for the areas below the frontier of need expected health care utilization. Although, the HCD profile measures inequity experienced only in this part of the areas, their measure of inequity takes into account the whole distribution of health care in all the areas. For all those areas above the frontier of health care utilization, the HCG vector returns a zero entry. Hence, the HCD profile is not able to measure inequity experienced by the non-deprived in health care areas. However, if some areas are above the frontier of need expected utilization, then at least one area will be deprived in health care (Proposition 1). Thus, the HCD profile will be indirectly able to measure this source of inequity in the population.

Finally, Proposition 2 states an unequivocal relationship between *the absence of deprivation* and *horizontal equity* in the whole population of areas. Therefore, the absence of deprivation in health care is a desirable condition since it guarantees horizontal equity for *all* the areas.

3. HCD-dominance and HCD-indices

In the inequality literature, the Lorenz dominance criterion requires a minimal set of value judgements³ in order to compare inequality in the distribution of income of different populations. This results in robust, but incomplete rankings. Inequality indices (e.g. the Gini, Theil or Atkinson index) play an important role whenever the Lorenz dominance criterion is not capable of ordering two distributions. These indices incorporate a larger number of value judgements, but are able to produce complete rankings. In this section, we follow the same axiomatic approach and illustrate the normative assumptions which characterize the dominance relation between HCD profiles and the class of indices consistent with HCD-dominance rankings.

Consider the class of indices Δ defined over the HCG vector, x , that satisfies the following normative properties:

1. Symmetry or anonymity axiom: $\Delta(x)$ is not influenced by permutations in the values of x .
2. Replication invariance axiom: if we can obtain x' from x by replication of the population, then $\Delta(x') = \Delta(x)$.
3. Strict monotonicity axiom: if x' is obtained from x by reducing some HCGs, then $\Delta(x') < \Delta(x)$.

³ These are the symmetry or anonymity axiom and the Pigou-Dalton principle of transfers.

4. Equality preferring axiom: if x' is obtained from x by mean preserving equalizations, then $\Delta(x') < \Delta(x)$. This means that if we transfer health care from some health care deprived area to some area that is more deprived, the value of $\Delta(\cdot)$ increases. The same result holds if we transfer health care from areas non-deprived to areas deprived in health care.
5. Modified focus axiom: if we obtain x' from x transferring health care across the non-deprived in health care, then $\Delta(x') = \Delta(x)$.

Then:

For any vector x' and x , x' HCD-dominates x if and only if $\Delta(x') < \Delta(x)$ for all $\Delta(\cdot) \in \Delta$. (Shorrocks, 1998; Jenkins and Lambert, 1997 and 1998)

Axioms 1 to 4 are widely accepted in the inequality literature and so we will not comment on these further (Lambert, 2001). Conversely, the focus axiom is chiefly used in the literature on poverty and has important implications for the inequity measure that we obtain. First, it implies that inequity among health care deprived areas should not be offset by inequity among health care non-deprived areas. Specifically, greater variance in the distribution of the gap of the former is not compensated by less variance in the distribution of the gap of the latter. Second, in ranking inequity between populations of areas *having equals average HCG* what should matter is inequity among the deprived in health care, not among the non-deprived in health care. This implies HCD ranks give zero weight to inequity to all those areas using more health care than need expected. However, this is not equivalent to excluding the top of the distribution of health care from the inequity measure that we obtain. For instance, if the utilization of health care non-deprived areas increases, while utilization of health care deprived areas remains unchanged, then the HCD profile will rise since the value of h^* for each observation is now slightly higher. Only transfers among health care non-deprived areas (so long as this does not result in a non-deprived area becoming deprived in health care) will not be captured by the HCD profile. This characteristic distinguishes the modified focus axiom defined above from the usual focus axiom delineated in the literature on poverty: an exogenous change in the utilization of health care non-deprived areas is captured by our class of indices.

Analysis of the distribution of health care across areas non-deprived in health care is achievable under this approach. We just need to invert the sign in the definition of the HCGs (i.e. $x_i = \max(h_i - h_i^*; 0)$) in order to capture the upper part of the health care distribution. The need-expected distribution of health care, h^* , is the same as that for the deprived in health care. Thus, we can easily define health care affluence profiles (HCAs) and obtain a measure of inequity as unequal access for equal need for the non-deprived in health care.

The HCD-dominance relation produces ranks that satisfy all the normative axioms listed above and are robust to any specification of the indices in the class Δ . This means that the HCD ranks are consistent with any social preferences for inequity and deprivation that satisfy the axioms. In particular, HCD-dominance ranks are consistent with preferences concerned exclusively with those areas having the largest HCG (Rawlsian preferences), and with preferences that make trade-offs between the largest and the average HCG in the population of areas. When the HCD profiles cross there is no dominance relation between distributions, and thus it is necessary to specify an index or a set of indices in the class $\Delta(\cdot)$ in order to obtain a complete ranking. This is equivalent to assuming some particular specification of social preferences. Complete rankings are obtained at the price of a loss of robustness, since they are no longer consistent with the entire preference spectrum defined above.

The Foster, Greer and Thorbecke (1984) class of poverty indices belong to the set $\Delta(\cdot)$ and are valuable candidates for the inequity analysis⁴. They can be defined as a function of HCGs and used as indices of horizontal inequity and deprivation in health care. Following the framework delineated in Section 2, we obtain:

$$\Delta_{\alpha}^{FGT}(x) = \int_0^{\xi} x^{\alpha} dF(x) \quad (13)$$

Where α is a parameter that defines the social preferences for the distribution of inequity among the areas deprived in health care. When $\alpha = 1$, the FGT-index equals μ and we obtain inequity ranks consistent with preferences that care only about the average HCG in the population of areas. When $\alpha > 1$, the transfer axiom holds and the inequity ranks are consistent with preferences that allow trade-offs between the distribution of the HCG and the value of μ . When $\alpha \rightarrow \infty$, no trade-off is allowed and inequity is ranked according to the area having the largest HCG in the population (Rawlsian preferences). A useful characteristic of the FGT family of indices is their additive decomposability property: if the population of areas is partitioned into K subgroups, then the overall value of the index can be expressed as a weighted sum of the value of the index in each subgroup. Therefore, the measure of inequity is subgroup consistent: a reduction in the inequity of one group will result in a reduction of inequity in the total population of areas. This property can be expressed as:

$$\Delta_{\alpha}^{FGT}(x) = \sum_{k=1}^K \pi_k \Delta_{\alpha}^{FGT}(x_k) \quad (14)$$

Where π_k is the proportion of areas in the subgroup k and x_k is the HCG vector for the group k .

4. Analysis of the GP fundholding reform

This section uses HCD profiles to measure the effect of GP fundholding on equity in the English NHS. The fundholding scheme was part of the ‘‘Internal Market’’ reforms implemented in the NHS during the years 1991-1999. One objective of the reform was to improve cost-efficiency and quality in the delivery of health care by creating local public payers responsible for purchasing care from local hospital providers and thereby introducing ‘‘quasi-market’’ hospital competition into the NHS. Two types of payer were introduced: geographically defined District Health Authorities and GP fundholders (GPFHs). GP fundholding was a voluntary scheme: the GPs who decided to become FHs received a budget to purchase a wide range of secondary care for the patients on their practice list. They could reinvest any savings in their practice, including improvements to premises and the range of services offered, both of which would potentially increase the capital value of the practice owned by the GP principals. Those GPs who did not join the scheme bore no cost in referring patients to hospital.

Many empirical studies have investigated the consequences of the scheme for secondary care admissions (Coulter and Bradlow, 1993; Propper, et al., 2002; Dusheiko et al., 2006). Virtually all these studies use a regression approach to evaluate the effect of fundholding and do not focus on the consequences for equity. The main contribution of our study is to analyse the effect of fundholding on equity, and to do so in a way consistent with a wide range of social preferences.

⁴ The FGT indices are not new to health economics. Madden (2006) has applied this class of indices in the analysis of the distribution of Body Mass Index scores in Ireland, but to our knowledge they have not been applied to equity in the delivery of health care.

4.1. Data

We use a small area panel dataset composed of over 8,000 electoral wards covering the whole of England from 1989 to 2001.

Data on hospital utilisation was obtained from the Hospital Episode Statistics (HES) for financial years 1989/90 through 2001/2. This includes data on all NHS hospital inpatient episodes in England, but excludes private hospital admissions. In our analysis we use data on all finished first consultant episodes for elective primary total prosthetic replacement of hip joint, excluding revisions, for all adults aged over 44 by age, sex and frozen 1991 ward code.

Data on the characteristics of ward resident populations in 1991 and 2001 was obtained from the respective National Censuses (Office of Population Censuses and Surveys 1991, Office of National Statistics 2001). Census 2001 data was allocated to frozen 1991 wards at the level of 2001 Census Output Areas, which each contain around 125 households. This was realized using geographical software (MapInfo), by matching each Output Area to the ward which contained the “population centroid” of that Output Area (i.e. the point of maximal population density). Census data was then linearly interpolated between 1991 and 2001 endpoints to estimate population characteristics in the years 1992-2000. Demographic and socio-economic characteristics in 1989 and 1990 were assumed to be unchanged with respect to the 1991 census point.

The Census-based Townsend deprivation Z-score is used as an indicator of socio-economic status (Carstairs, 2000).

The ward population size aged over 44, split by five-year bands (from 44 up to over 90) and by sex is used as indicator of need. Few elective hip procedures are performed on patients under 45 (Dixon *et al.*, 2004).

We compute an index of GP fundholding penetration in order to capture the effect of the reform on secondary care admissions. The index is time varying and is defined as the proportion of the ward population registered with a fundholding practice. The sources of data for this index includes Prescription Pricing Authority data from 1998/9 on all practices in England including practice code, fundholding type and start/end date; Attribution Data Set data on practice registrations for 1999 2000; and record linkage data from the Codes Service Database 2001.

4.2 Methods of analysis

Elective hip replacements for all adults aged over 44 are used as an indicator of health care utilization. The vector of need-expected utilization, h^* , is computed following Equations (3) and (4):

$$h_i^* = \sum_{j=1}^J p_{ji} h_j \quad (3)$$

Where $J = 10$ are need classes (five age classes for men and five for women aged from 44 up to over 90); p_{ji} is the number of people in the ward i in the age and sex stratum j ; h_j is the national utilization rate for age and sex stratum j , i.e. the national count for England in the current year divided by the national population in that year. This specification of h^* facilitates inequity comparison between different years since it allows for the increase in national utilization over the period considered.

The HCG vector defined in section 1 is obtained as: $x_i = \max(h_i^* - h_i; 0)$, where h_i is the observed count for elective hip replacements in ward i . We account for potential bias due to heterogeneity in the ward population size using indirect standardization (O'Donnell et al., 2008):

$$x_i^{is} = x_i - \hat{x}_i + \bar{x} \quad (15)$$

$$\hat{x}_i = \hat{\beta}_0 + \hat{\beta}_1 p_i + \hat{\beta}_2 \bar{d} + \hat{\beta}_3 \bar{g} \quad (16)$$

The HCGs, x_i , are regressed against population size p_i controlling for level of deprivation d_i and fundholding penetration g_i . Equation (15) gives the indirectly standardized HCGs at the average population size effect. This variable can be interpreted as the HCGs that we would observe if the population in each ward was equal to the average population⁵.

Trends in inequity over time were tracked for all the health care deprived wards. The FGT indices are computed for the whole population of English wards from 1989 to 1991 and decomposed by level of *socio-economic deprivation*, as measured by the Townsend index, in four quartiles (most affluent 1st, most affluent 2nd, most affluent 3rd, most deprived quartile). The parameter α is set: $\alpha = 1$, producing ranks consistent with social preferences that care only for the average HCG; $\alpha = 2$, producing ranks consistent with social preferences that trade-off inequality in the distribution of HCGs with the magnitude of the average HCG. Robust standard errors are computed accounting for heterogeneity in the ward population size using statistics developed for the index in the literature on poverty⁶ (Kakwani, 1993; Joliffe et al., 2004).

The impact of the GP fundholding reform on equity is analysed by focusing on the most socio-economically deprived quartile of electoral wards deprived in health care. We expect that, if the reform has an impact on equity this is more likely to be evident in those wards where inequity has consistently been present over the period of time considered (Figure 2 and Figure 3). Also, selecting the most socio-economically deprived quartile of wards has two advantages. First, we achieve some additional control for unobserved heterogeneity in the need for hip replacement across the wards: if unobservable need for hip replacement is positively correlated with the level of deprivation of the wards, selecting the most socio-economically deprived wards would reduce such heterogeneity in the analysis. Second, we exclude that part of the population that is more likely to select out from the sample by opting for private health care treatment, which is not observable in our dataset. This is a particularly relevant issue for hip replacement, since one quarter of all the procedures recorded in England during the period examined were performed by the private sector (Williams and Rossiter, 2004). Thus, potential bias coming from sample selection should be considerably reduced.

The most socio-economically deprived quartile of wards is initially partitioned into two groups: the control group composed of wards having a proportion of people registered in a FH practice lower than 20%, and the treatment group having a proportion of people registered in a FH practice larger than 20%. In order to check the robustness of our findings further subgroups are considered, dividing the treated group according to increasing fundholding penetration⁷ (20-40%, 40-60%, more than 60%, 40-100%). We compute HCD profiles for each group from 1991 to 1996 and test for

⁵ Also, we attempt to control for unobservable heterogeneity of the HCGs in the need for hip replacements that is not captured by demographic variables. The standardized illness ratio defined in section 4.1 is used as a proxy of the need for hip replacements. However, we find that the latter is not significant in explaining the variation in the HCGs after controlling for the ward population aged over 44.

⁶ These statistics are validated using bootstrapped standard errors.

⁷ Further stages of decomposition are not achievable for 1991 and partially achieved for 1992 wards, because of insufficient observations having fundholding penetration higher than 40% in those early years (only 58 wards in 1991 and 130 wards in 1992).

HCD-dominance implementing the statistical test developed by Davidson and Duclos (2000). The HCD profiles are displayed in Figures 4 and the results of the tests are in Table 2.

FGT indices (parameters $\alpha=1$ and $\alpha=2$) are computed for the groups of wards defined above from 1991 to 1996. An index of relative inequity is computed dividing within-group FGT indices by the FGT index for the sum of the groups:

$$r_{\alpha}^{FGT}(x_k) = \frac{\Delta_{\alpha}^{FGT}(x_k)}{\Delta_{\alpha}^{FGT}(x_K)} = \frac{\Delta_{\alpha}^{FGT}(x_k)}{\sum_{k=1}^K \pi_k \Delta_{\alpha}^{FGT}(x_k)}, \forall k \in K \subseteq N \quad (17)$$

K is the number of subgroups of wards defined by increasing degree of fundholding penetration in the most socio-economically deprived quartile of wards (i.e. a subset of the total number of wards N). “The relative FGT index” shows the inequity share of each group according to the level of fundholding penetration: a ratio above 1 indicates more inequity than the average, i.e. the inequity measured in the sum of the groups. Results are shown in Table 3 and in Figure 5 and Figure 6. We test for the hypothesis of no significant difference in the share of inequity between all groups from 1991 to 1996 using bootstrapped standard errors.

The test for the hypothesis of no treatment effect is implemented using a difference-in-differences approach. First, we compute the difference between treated groups before and after the fundholding reform was implemented, then we subtract from this value the difference between the control group before and after the reform. 1989/90 was set as benchmark financial year, since it falls two years before the start of the FH scheme, allowing us to account for potential strategic behaviour of GPs preparing to become FHs in 1991 (Croxson, et al., 2001). This difference-in-differences approach can be formulated as:

$$DID = (\Delta_{f,t}^{FGT} - \Delta_{f,'89}^{FGT}) - (\Delta_{nf,t}^{FGT} - \Delta_{nf,'89}^{FGT}) \quad (18)$$

where f is the treated group and nf is the control group. The DID allows us to control for time confounders and for self-selection. The DID identifies the treatment effect under the assumption that the effect of time and policy confounders follow a common time trend for fundholding and non fundholding practices (Jones, 2007). The results are reported in Table 4.

4.3 Results

HCD profiles and FGT indices can be interpreted both as measures of deprivation in the access to elective hip replacements and as measures of inequity with respect to those wards which result to be deprived in health care. We will follow the latter interpretation in explaining the results of our analysis.

Figures 2 and 3 show that among those wards deprived in health care, the most socio-economically deprived quartile experiences the greatest inequity (or equivalently the greatest deprivation in health care) over all periods considered. The remaining quartiles have no significant differences in their values of the FGT indices. This result is robust to two alternative specifications of the social preference for the distribution of inequity captured by the parameters $\alpha=1$ and $\alpha=2$ of the index. Our findings are consistent with a large part of the literature on socio-economic inequalities and inequity in elective hip replacement in the NHS (Chaturvedi and Ben-Shlomo, 1995; Cookson et al., 2007). The difference between the average HCG of the most socio-economically deprived quartile and the average HCG of other quartiles ranges from 0.66 elective hip replacements in 1989 to 1.54 in 1993

(Figure 2). This is a quite substantial disparity if we consider that the (population-over-44 weighted) average number of elective hip replacements for each ward was 3.57 in 1989 and 4.86 in 1993.

Table 2. HCD-dominance ranks in the most socio-economically deprived quartile of wards; all observations are grouped by level of fundholding penetration (0-100%); test for dominance from Davidson and Duclos (2000).

| Population subgroups | Years | | | | | |
|----------------------|-------|--------|--------|--------|--------|--------|
| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 |
| A vs B | SSD** | SSD** | no dom | no dom | no dom | no dom |
| A vs C | - | no dom | no dom | no dom | no dom | no dom |
| A vs D | - | - | SSD** | no dom | no dom | no dom |
| A vs E | - | - | SSD** | no dom | SSD** | SSD* |
| A vs F | - | SSD** | SSD** | no dom | no dom | no dom |
| C vs D | - | - | no dom | no dom | no dom | no dom |
| C vs E | - | - | SSD** | no dom | SSD* | SSD** |
| C vs F | - | no dom | SSD** | no dom | no dom | no dom |
| D vs E | - | - | no dom | no dom | no dom | no dom |

Note: A = 0 to 20% of people registered in a GPFH practice; B = 20% to 100%;
C = 20% to 40%; D = 40% to 60%; E = 60% to 100%; F = 40% to 100%.
SSD** = second order stochastic dominance significant at 5%;
SSD* = significant at 10%; no dom = not significant.

In the most socio-economically deprived quartile of wards, higher FH penetration is associated with less inequity for all those wards deprived in health care (Table 2, Figure 4). There is a clear dominance rank by increasing degree of FH penetration in 1991 to 1993. However, this dominance relationship holds only with respect to the top FH penetration group (i.e. 60-100% FH penetration) in 1995 and 1996, and is not statistically significant in 1994. Turning to the FGH indices, inequity falls with increasing degree of FH penetration over all the years of the reform (Table 3, Figures 5 and 6). In 1993, people in socio-economically deprived wards with over 60% FH penetration have an average HCG 29% smaller than the average for wards in the same most socio-economically deprived quartile of electoral wards (Table 3, Figure 5). The positive relationship between FH penetration and equity reduces in intensity after 1993 and holds only for the top FH penetration groups, confirming the dominance ranks in Table 2. We can interpret this change in the trend as a break in the characteristics of the FHs: early waves of FHs were more responsive to equity in the distribution of access to elective hip replacements, while late waves of FHs were less responsive. Previous studies support the hypothesis of a break in the characteristics of the GPFHs that joined the scheme after 1993 (Baines and Whyntes, 1996). The authors bring evidence that FHs in the first

three waves were more likely to meet a set of quality and efficiency targets than the FHs in subsequent waves. An alternative hypothesis is that the change in the trend is a consequence of a change in the behaviour of the early wave FHs. However, neither the former nor the latter hypothesis can be tested in our study. Since our unit of analysis is the electoral ward, we are not able to adequately disentangle the effects of individual waves of FHs.

Table 3. Differences in inequity share in most socio-economically deprived quartile of electoral wards of England; all observations are grouped by level of fundholding penetration (0-100%); relative FGT indices; bootstrapped standard errors.

| Population subgroups | Preference parameter | Years | | | | | |
|----------------------|----------------------|---------|---------|---------|---------|---------|---------|
| | | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 |
| A – B | $\alpha = 1$ | 16.8%** | 19.2%** | 14.5%** | 8.8%** | 4.3% | 1.6% |
| | $\alpha = 2$ | 21.9%** | 22.9%** | 14.8%** | 9.0% | 4.4% | -3.0% |
| A – C | $\alpha = 1$ | - | 18.5%** | 8.6% | 4.9% | -0.2% | -4.0% |
| | $\alpha = 2$ | - | 21.2%** | 6.8% | 2.6% | -0.9% | -16.1%* |
| A – D | $\alpha = 1$ | - | - | 21.6%** | 13.3%** | 8.8% | 2.2% |
| | $\alpha = 2$ | - | - | 23.3%** | 16.9%* | 6.5% | 1.5% |
| A – E | $\alpha = 1$ | - | - | 29.1%** | 13.6%** | 9.8%* | 10.3%* |
| | $\alpha = 2$ | - | - | 36.4%** | 15.8%* | 15.7%** | 13.2%* |
| A – F | $\alpha = 1$ | - | 20.5%** | 23.9%** | 13.4%** | 9.2%* | 5.7% |
| | $\alpha = 2$ | - | 26.5%** | 27.3%** | 16.5%** | 10.3% | 6.6% |
| C – D | $\alpha = 1$ | - | - | 13.1%* | 8.4% | 9.0% | 6.3% |
| | $\alpha = 2$ | - | - | 16.4% | 14.3% | 7.4% | 17.6%* |
| C – E | $\alpha = 1$ | - | - | 20.5%** | 8.8% | 9.9%* | 14.3%** |
| | $\alpha = 2$ | - | - | 29.5%** | 13.3% | 16.6%** | 29.3%** |
| C – F | $\alpha = 1$ | - | 2.0% | 15.3%** | 8.6% | 9.4%* | 9.8%* |
| | $\alpha = 2$ | - | 5.3%* | 20.4%** | 13.9% | 11.2% | 22.7%** |
| D – E | $\alpha = 1$ | - | - | 7.4% | 0.3% | 0.9% | 8.0% |
| | $\alpha = 2$ | - | - | 13.1% | -1.0% | 9.2% | 11.7% |

Note: A = 0 to 20% of people registered in a GPFH practice; B = 20% to 100%;
C = 20% to 40%; D = 40% to 60%; E = 60% to 100%; F = 40% to 100%.
** = significant at 5%; * = significant at 10%.

The difference-in-differences analysis shows that a large part of the observed equity associated with FH penetration is due to self-selection. In 1991, among the most socio-economically deprived quartile of wards, more equitable practices tended to self-select into the scheme (Table 4). In 1992 and 1993, there is evidence of a positive treatment effect of the FH reform on equity in the distribution of hip replacements. The magnitude of the effect is quite substantial: in 1993 the average HCG in the top FH penetration group reduces by 1.12 elective hip replacements with respect to the population weighted average number of elective hip replacements of 4.86 for each ward. However, we are not able to identify to what extent the effect is explained by the individual waves of FHs that join the scheme in 1991, 1992 or in 1993, since we are not able to control for FH waves. The treatment effect is not statistically significant after 1993. This finding is consistent with evidence that early wave FHs were more "enthusiastic" and "active" than late wave FHs (Le Grand et al., 1998). It suggests that the former may have been willing to increase hospital referral rates for

severe osteoarthritis among "high-need" individuals living in socio-economically deprived wards, relative to the rest of the population. However, other possible explanations cannot be ruled out in the absence of practice level data that could disentangle selection and treatment effects by FH waves and in the absence of data on GP referral rates as opposed to hospital admission rates.

Table 4. Difference in differences estimates from FGT indices; population subgroups defined by level of fundholding penetration (0-100%); non-treated group: 0-20% FH penetration; reference year: 1989.

| Treated groups | Preference parameter | Years | | | | | |
|----------------|----------------------|--------|----------|----------|--------|--------|--------|
| | | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 |
| B | $\alpha = 1$ | -0.285 | -0.461** | -0.620** | -0.222 | 0.062 | 0.226 |
| | $\alpha = 2$ | -2.02 | -2.985* | -3.9** | -1.193 | 0.52 | 2.710* |
| C | $\alpha = 1$ | - | -0.304 | -0.493** | -0.267 | 0.100 | 0.116 |
| | $\alpha = 2$ | - | -2.069 | -2.958 | -1.37 | 1.072 | 2.928 |
| D | $\alpha = 1$ | - | - | -0.689** | -0.101 | -0.063 | 0.335 |
| | $\alpha = 2$ | - | - | -4.304* | -0.704 | 0.059 | 2.637 |
| E | $\alpha = 1$ | - | - | -1.117** | -0.279 | 0.143 | 0.270 |
| | $\alpha = 2$ | - | - | -7.827** | -1.423 | -0.278 | 2.470* |
| F | $\alpha = 1$ | - | -0.788** | -0.819** | -0.168 | 0.022 | 0.306 |
| | $\alpha = 2$ | - | -4.891* | -5.379** | -0.973 | -0.085 | 2.559* |

Note: A = 0 to 20% of people registered in a GPFH practice; B = 20% to 100%;
 C = 20% to 40%; D = 40% to 60%; E = 60% to 100%; F = 40% to 100%.
 ** = significant at 5%; * = significant at 10%.

4.4 Robustness checks

Please, refer to the original version of the paper:
<http://www.york.ac.uk/res/herc/research/hedg/wp.htm>

5. Conclusions

The methodological contribution of this paper is to introduce some developments in the literature of poverty and deprivation to the analysis of inequality and inequity in health care. The HCD profiles define deprivation in health care as receiving less care than needed. This definition is strictly related to inequity, since a common interpretation of equity in health care is that resources have to be allocated with respect to individuals' needs rather than individuals' preferences. Thus, if some minorities in the population have inadequate access with respect to their need, this is generally regarded as an issue of social justice. In this context, the definition and the measurement of need for health care assumes a crucial role. The HCD profiles do not depend on a specific definition of need. In this paper, we have proposed a function of the average utilization observed in the total population as need estimator, following the tradition of the literature on health care inequality. In turn, this has led us to determine a link between horizontal equity and deprivation. However, alternative definitions of need can be specified, and thus alternative view on equity can be included in the HCD profiles.

A wide class of indices consistent with the HCD-dominance ranks can be introduced from the literature on poverty to provide measures of inequity and deprivation in health care. The FGT indices are members of this class and have some appealing properties. First, these indices are additive decomposable: inequity measured in the total population is a weighted average of inequity in the population subgroups. Hence, the measure of inequity is subgroup consistent: a reduction in the inequity of one group will result in a reduction of inequity in the total population. This property enables us to calculate different subgroups' contributions to overall inequity and to identify those groups suffering the largest inequity in the population. Often, the latter represent valuable information for the policy maker. Second, the FGT indices permit an explicit specification of social preferences for inequity and deprivation into a single parameter. Therefore, it is possible to test the robustness of the inequity ranks to alternative views about equity which endorse different trade-offs between the average HCG and inequality in the distribution of the HCGs.

The HCD profiles disregard information on inequity of one part of the distribution, i.e. those units having more utilization than their need-expected level. On one hand, focusing sharply in this way results in a graphical representation of useful information on the inequity experienced by those who are below their need expected utilization (i.e. the incidence the intensity and the inequality in the distribution of the HCGs). On the other hand, the HCD profiles represent inequity experienced only by one part of the population. However, we have shown in Section 3 that this measure of inequity takes into account information on the whole distribution of health care in the population of areas. Furthermore, it is possible to reverse the focus and look only at that part of the distribution receiving more than their need expected level of health care - i.e. an analysis of health care affluence profiles.

Section 4 illustrates an empirical application of the HCD profiles to measure the effect of the GP fundholding reform in the English NHS. Inequity in the distribution of elective hip replacements is measured as unequal treatment for equal need among those electoral wards deprived in health care. Inequity trends over time are analysed across socio-economic quartiles of wards. Measures of inequity are decomposed according to increasing level of FH penetration in the most socio-economically deprived quartile of wards. We produce HCD-dominance ranks associated with FH penetration, which are consistent with a large range of social preferences. The effect of the FHs reform on equity is identified using a difference-in-differences approach under two alternative specification of social preference for inequity embedded in the parameter α of the FGT index. The additive decomposability property of the FGT indices is crucial for the implementation of this estimation method, since it permits a decomposition of inequity measures by treated and non treated groups.

Our results show that in 1991 the lower level of inequity associated with fundholding is due to self-selection of the most equitable practices into the scheme. There is evidence of self-selection and also a positive treatment effect on equity in 1992 and 1993. The treatment effect is considerable. In 1993/4, the average health care gap between the actual and need-expected number of elective hip replacements in the top fundholding penetration wards fell by 1.12 hip replacements, compared to a population-weighted average number of elective hip replacement of 4.86 for each ward. However, we are not able to identify such effect by waves of FHs. Finally, in 1994 to 1996 the positive effect on equity is explained only by self-selection and is significant only in the group with the highest penetration of fundholding.

References

-Baines DL, Whyne DK. Selection bias in GP fundholding. *Health Economics* 1996;5: 139-140.

-Breslow NE, Day NE. *Statistical Methods in Cancer Research: The Design and Analysis of Cohort Studies*, vol.2. International Agency for Research on Cancer: Lyon; 1987.

- Chaturvedi N, Ben-Shlomo Y. From the surgery to the surgeon: does deprivation influence consultation and operation rates? *British Journal of Medical Practice* 1995;45; 127-131.
- Clayton D, Hills M. *Statistical Models in Epidemiology*. Oxford University Press: Oxford; 1993.
- Cookson R, Dusheiko M, Hardman G. Socio economic inequalities in small area use of elective total hip replacement in the English National Health Service in 1991 and 2001. *Journal of Health Services Research and Policy* 2007;12 Suppl 1, S1-10-7.
- Coulter A, Bradlow J. Effect of NHS reforms on general practitioners' referral patterns. *British Medical Journal* 1993;306; 433-437.
- Croxson B, Propper C, Perkins A. Do doctors respond to financial incentives? UK family doctors and the GP fundholder scheme. *Journal of Public Economics* 2001;79; 375-398.
- Davidson R, Duclos J-Y. Statistical inference for stochastic dominance and for the measurement of poverty and inequality. *Econometrica* 2000;68; 1435-1464.
- Dixon T, Show M, Ebrahim S, Dieppe P. Trends in hip and knee joint replacement: socioeconomic inequalities and projections of need. *Annals of the Rheumatic Diseases* 2004;63; 825-830.
- Dusheiko M, Gravelle H, Jacobs R, Smith P. The effect of financial incentives on gatekeeping doctors: evidence from a natural experiment *Journal of Health Economics* 2006;25; 449-478.
- Foster J, Greer J, Thorbecke E. A class of decomposable poverty measures. *Econometrica* 1984;52; 761-766.
- Gravelle H, Morris S, Sutton M 2006. Economic studies of equity in the consumption of health care. In: Jones AM (Eds), *The Elgar Companion to Health Economics*. Edward Elgar Publishing Limited: Cheltenham; 2006. p. 193-204.
- Jenkins SP, Lambert PJ. Three I's of poverty curves, with an analysis of UK poverty trends. *Oxford Economic Papers* 1997;49; 317-327.
- Jenkins SP, Lambert PJ 1998. Three 'I's for poverty curves and poverty dominance: TIPs for poverty analysis. In: Slotte DJ (Eds), *Research on Economic Inequality*. Jay Press Inc.: London; 1998. p. 39-56.
- Joliffe D, Datt G, Sharma M. Robust Poverty and Inequality Measurement in Egypt: Correcting for Spatial-price Variation and Sample Design Effects. *Review of Development Economics* 2004;8; 557-572.
- Jones AM. Panel data methods and applications to health economics. HEDG working Paper 07/18, University of York, 2007; <http://www.york.ac.uk/res/herc/research/hedg/wp.htm>.
- Kakwani N. Statistical inference in the measurement of poverty. *Review of Economics and Statistics* 1993;75; 632-639.
- Lambert P. *The Distribution and Redistribution of Income*. Manchester University Press: Manchester; 2001.
- Le Grand J, Mays N, Mulligan K. *Learning from the NHS internal market: a review of the evidence*. King's Fund Publishing: London; 1998.
- Madden D. Body mass index and the measurement of obesity. HEDG working paper 06/11, University of York, 2006; <http://www.york.ac.uk/res/herc/research/hedg/wp.htm>.
- Morris S, Sutton M, Gravelle H. Inequity and inequality in the use of health care in England: an empirical investigation. *Social Science & Medicine* 2005;60; 1215-1266.
- O'Donnell O, van Doorslaer E, Wagstaff A, Lindelow M. *Analyzing Health Equity Using Household Survey Data*. World Bank Publications: Washington DC; 2008.
- Propper C, Croxson B, Shearer A. Waiting times for hospital admissions: the impact of GP fundholding. *Journal of Health Economics* 2002;21; 227-252.
- Shorrocks A 1998. Deprivation profiles and deprivation indices. In: Jenkins SP, Kapteyn A, Van Praag BMS (Eds), *The Distribution of Welfare and Household Production: International Perspectives*. Cambridge University Press: London; 1998. p. 250-267.
- van Doorslaer E, Wagstaff A. Equity in the delivery of health care: some international comparisons. *Journal of Health Economics* 1992;11; 389-411.
- van Doorslaer E, Koolman X, Jones AM. Explaining income-related inequalities in doctor utilization in Europe. *Health Economics* 2004;13; 629-647.
- van Doorslaer E, Messaria C, Koolman X. Inequalities in access to medical care by income in developed countries. *Canadian Medical Association Journal* 2006;174; 177-183.
- Wagstaff A, van Doorslaer E, Paci P. On the measurement of horizontal inequity in the delivery of health care. *Journal of Health Economics* 1991;10; 169-205.
- Wagstaff A, van Doorslaer E. Measuring and testing for inequity in the delivery of health care. *Journal of Human Resources* 2000a;35; 716-733.
- Wagstaff A, van Doorslaer E 2000b. Equity in health care finance and delivery. In: Culyer AJ, Newhouse JP (Eds), *Handbook of Health Economics*. Elsevier Science BV: Amsterdam; 2000b. p. 1804-1862.
- Williams J, Rossiter A. *Choice: the evidence*. Social Market Foundation: London; 2004.

Figure 2.




Figure 3.




Figure 4. HCD profiles by presence of FHs in most socio-economically deprived quartile of electoral wards.




Figure 5. Relative FGT index decomposed by presence of FHs (average inequity in the total population = 1; social preference parameter $\alpha = 1$).

